

Table 1: Controlled clinical trials of fermented wheat germ extract (FWGE) for cancer

Source: CAM-Cancer Consortium. Fermented wheat germ extract [online document]. March, 2025.

Study	Methods	Participants	Interventions	Results	Comments
Randomized controlled trials					
Demidov 2008	2-arm, randomized, open label, parallel	58 patients with malignant melanoma (stage III)	Darcarbazine with or without FWGE	Improved survival rates and progression free survival after seven years of follow up in the FWGE group	Small study with high risk of selection bias
Prospective, non-randomized controlled trials					
Jakab 2003	2-arm, prospective, nonrandomized, open-label, parallel	176 patients with colorectal cancer	Oncological therapy (radio- and/or chemotherapy) with or without FWGE	Reduced metastases, higher overall survival rates and longer progression free survival in the FWGE group	Study with high risk of selection bias
Sukkar 2008	2-arm, prospective, nonrandomized, open label, parallel.	55 patients diagnosed with head and neck cancer (stage IIIa, IIIb and IV)	Oncological therapy (surgery, radio- and/or chemotherapy) with or without FWGE	Significantly better quality of life in the FWGE group	Small study with high risk of selection and detection bias low reporting quality
Garami 2004	2-arm, prospective, nonrandomized, controlled, open label, parallel.	22 children with various tumours	Different chemotherapy regimen with or without FWGE	Significantly reduced incidence of neutropenic fever in the FWGE group	Low reporting quality. Small study with high risk of selection bias.
Barabàs 2006	2-arm, prospective, parallel	45 patients diagnosed with head and neck cancer (UICC stage II and III)	Oncological therapy (surgery, radio- and/or chemotherapy) with or without FWGE	Significantly fewer local relapses and fewer progression events in the FWGE group	Small study with high risk of selection and detection bias. Low reporting quality.